

Collisional cooling of ultracold molecules

<https://doi.org/10.1038/s41586-020-2141-z>

Hyungmok Son^{1,2✉}, Juliana J. Park¹, Wolfgang Ketterle¹ & Alan O. Jamison^{1,3}

Received: 22 July 2019

Accepted: 27 January 2020

Published online: 8 April 2020

 Check for updates

Since the original work on Bose–Einstein condensation^{1,2}, the use of quantum degenerate gases of atoms has enabled the quantum emulation of important systems in condensed matter and nuclear physics, as well as the study of many-body states that have no analogue in other fields of physics³. Ultracold molecules in the micro- and nanokelvin regimes are expected to bring powerful capabilities to quantum emulation⁴ and quantum computing⁵, owing to their rich internal degrees of freedom compared to atoms, and to facilitate precision measurement and the study of quantum chemistry⁶. Quantum gases of ultracold atoms can be created using collision-based cooling schemes such as evaporative cooling, but thermalization and collisional cooling have not yet been realized for ultracold molecules. Other techniques, such as the use of supersonic jets and cryogenic buffer gases, have reached temperatures limited to above 10 millikelvin^{7,8}. Here we show cooling of NaLi molecules to micro- and nanokelvin temperatures through collisions with ultracold Na atoms, with both molecules and atoms prepared in their stretched hyperfine spin states. We find a lower bound on the ratio of elastic to inelastic molecule–atom collisions that is greater than 50—large enough to support sustained collisional cooling. By employing two stages of evaporation, we increase the phase-space density of the molecules by a factor of 20, achieving temperatures as low as 220 nanokelvin. The favourable collisional properties of the Na–NaLi system could enable the creation of deeply quantum degenerate dipolar molecules and raises the possibility of using stretched spin states in the cooling of other molecules.

The full potential of ultracold atoms was not realized until the advent of collision-based cooling methods such as evaporative and sympathetic cooling. Although atomic systems have been recently used to demonstrate laser cooling to quantum degeneracy, these schemes still require collisional thermalization^{9,10}. Therefore, many efforts have been made over the past 15 years¹¹ to achieve collisional cooling of ultracold molecules. Buffer gas cooling⁸ cannot achieve temperatures below 100 mK owing to the rapidly diminishing vapour pressure of buffer gases at such temperatures. Supersonic expansion⁷ can produce temperatures around 100 mK. Controlled collisions in crossed molecular beams¹² can decrease the laboratory-frame velocity of particles while narrowing the velocity distribution. However, this technique has not been demonstrated below about 500 mK. Merged supersonic beams can be used to study collisions at energies equivalent to a temperature of 10 mK (ref. ¹³).

Cooling below 100 mK calls for trapping molecules in magnetic or electrostatic traps and for good collisional properties (that is, a ratio of elastic to inelastic collisions much greater than 1). Such traps require preparing molecules in weak-field seeking states, which are never the absolute ground state, allowing inelastic state-changing collisions to eject the cold molecules from the trap. A variety of systems have been proposed for evaporative or sympathetic cooling of molecules^{11,14–18}. So far, elastic collisions have been observed clearly in O₂ at temperatures below 1 K (ref. ¹⁹) and possibly in OH radicals around 10 mK (ref. ²⁰

corrects an earlier report²¹). In the O₂ case, inelastic collisions prevent thermalization and collisional cooling.

In recent years, the assembly of molecules from ultracold atoms²² and the direct laser cooling of molecules^{23–26} have both expanded to new molecules and temperature regimes. These techniques have achieved molecular systems at temperatures less than 100 mK, raising the challenge of collisional cooling in the micro- and nanokelvin regimes. Optical traps enable trapping of the absolute ground state, which removes the concern of state-changing collisions. However, collisional cooling in the absolute ground state of chemically stable systems has not yet been realized²⁷. By contrast, chemically stable molecular species have shown anomalously high inelastic loss rates that preclude collisional cooling, possibly owing to collision complex formation²⁸ or interactions with optical trapping beams²⁹.

In this study we observe sympathetic cooling of rovibrational ground-state triplet ²³Na⁶Li molecules by ²³Na atoms, both of which are prepared in their upper stretched hyperfine spin states (that is, states with both nuclear and electronic spins aligned along the direction of the magnetic bias field). Although sympathetic cooling of one atomic species by another has been observed in various ultracold atomic mixtures³⁰, triplet NaLi was considered unlikely to have sufficiently good collisional properties to support such cooling. NaLi has energetically allowed chemical reactions, even in the electronic ground state (that is, a singlet state), and the triplet state has an electronic excitation energy of 0.8 eV or 10,000 K.

¹Research Laboratory of Electronics, MIT-Harvard Center for Ultracold Atoms, Department of Physics, Massachusetts Institute of Technology, Cambridge, MA, USA. ²Present address: Department of Physics, Harvard University, Cambridge, MA, USA. ³Present address: Department of Physics and Astronomy, Institute for Quantum Computing, University of Waterloo, Waterloo, Ontario, Canada. ✉e-mail: hson@g.harvard.edu




Fig. 1 | Experimental setup. The Na atoms (yellow circles) and NaLi molecules (yellow and red circles on black sticks) are trapped in a 1D optical lattice formed by a 1,596-nm laser, which is retro-reflected. The magnetic field, which defines the quantization axis, is coaxial with the lattice beam. The free atoms that remain after the formation of the ground-state molecules are removed by resonant light in the radial direction (y axis in the figure). Stimulated Raman adiabatic passage (STIRAP) is performed with circularly polarized beams of two wavelengths (833 nm and 819 nm) that propagate along the axial direction (z axis in the figure). σ^+ and σ^- represent left-handed and right-handed circular polarization, respectively. The ground-state molecules are detected by absorption imaging along the axial direction of the atoms resulting from the dissociation of NaLi.

Furthermore, theoretical studies on various systems have explored the possibility of suppressing inelastic collisions and reactions by spin polarization, giving pessimistic predictions for triplet molecules and more favourable ones for doublet molecules^{17,31–34}. Nonetheless, here we report clear thermalization and sympathetic cooling of triplet NaLi to 220 nK. We observe 20-fold increases of phase-space density (PSD), opening the possibility of collisional cooling to deep quantum degeneracy.

Our experimental setup is summarized in Fig. 1. Similarly to our previous work^{35,36}, we produce about 3.0×10^4 NaLi molecules in the rovibrational ground state of the triplet potential using a mixture of Na and Li atoms (further details in Methods). We also prepare about 1.0×10^5 Na atoms in the upper stretched hyperfine state (in the low-field basis, $|F, m_f\rangle = |2, 2\rangle$, where F and m_f are the hyperfine and magnetic quantum numbers, respectively) in a one-dimensional (1D) optical lattice formed by 1,596-nm light. Owing to the differential polarizability at 1,596 nm, molecules feel a deeper trapping potential than atoms. This results in a sudden increase of the potential energy as atoms associate and form molecules (see Fig. 2a). Immediately after production, the effective temperature of the molecules is $2.80(6)$ μK and the temperature of Na atoms is $2.42(3)$ μK (all uncertainties are as defined in the legend of Fig. 2). As the molecules thermalize with the Na atoms and a hot fraction of atoms evaporates out of the trap, the temperatures of both particles settle to $2.23(6)$ μK (see Methods for molecular thermometry).

Although this initial settling of temperatures hints at sympathetic cooling, we are able to see much stronger effects by forced cooling and heating of Na atoms. We evaporate Na atoms with almost no loss of molecules by taking advantage of the particles' different polarizabilities: $\alpha_{\text{NaLi}}/\alpha_{\text{Na}} = (m\omega^2)_{\text{NaLi}}/(m\omega^2)_{\text{Na}} \approx 2.6$, where α_i is the polarizability of particle i , ω is the angular frequency for oscillation in the trap and m is the particle mass. The curve in Fig. 2c shows the thermalization between molecules and atoms with a large initial temperature difference after an exponential evaporative cooling ramp followed by a recompression of the trap (see Fig. 2 legend). As we hold the particles in the trap, their temperatures approach each other. Owing to the large particle number ratio, $N_{\text{Na}}/N_{\text{NaLi}} \approx 7$, the molecule temperature decreases by $0.68(9)$ μK after thermalization, whereas the temperature of Na atoms increases only by $70(30)$ nK. However, if the Na is removed immediately before the hold time, the molecule temperature remains fixed during the same period.




Fig. 2 | Thermalization of Na and NaLi. **a**, The trapping potential of molecules (red solid line) is deeper than that of Na atoms (black dashed line). This allows us to evaporate Na atoms with negligible loss of molecules. **b, c**, The molecule number (**b**) and temperature (**c**) are measured at various hold times after a 100-ms-long exponential evaporative ramp to a trap power of 0.21 W (7 μK trap depth), followed by a 10-ms-long recompression to 1.2 W (40 μK trap depth; trap-depth values are for NaLi throughout). In the number plot (**b**) the red dashed line is an exponential loss fit and the blue solid line is a two-body loss fit (details in the main text). The dashed line in the temperature plot (**c**) is an exponential fit. No temperature drop occurs in the absence of Na atoms. The exponential fit of the molecule temperature agrees well with the measurement but should be regarded as an interpolation to determine the initial slope of the thermalization. For this measurement, the Na number was about 1.5×10^5 . Data values represent the average and error bars represent the standard error of the mean, estimated from the fitting and statistical errors of 3–8 measurements.

As further evidence of thermalization, the sympathetic heating of molecules with hot atoms is shown in Fig. 3. We first prepare the atom–molecule mixture fully thermalized at 1.5 μK , after a 10-ms-long exponential evaporative ramp, followed by a 200-ms-long recompression of the trap to the initial trap power of 1.5 W (50 μK trap depth). Then, we selectively heat the atoms to 2.07(5) μK by sinusoidally modulating the trap amplitude at twice the sodium trap frequency ($2\omega_{\text{Na}} = 2\pi \times 920$ Hz) with a modulation amplitude of 20% of the trap depth for 100 ms. After




Fig. 3 | Sympathetic heating. After forced heating of Na atoms (see main text for details), the NaLi molecule temperature (red squares) rises and reaches that of Na atoms (black asterisk) as both particles thermalize. The temperature of molecules without hot atoms (blue circles) remains low. Values and error bars are estimated as in Fig. 2 from four measurements.




Fig. 4 | Evaporation sequences. The initial power of the 1,596-nm trapping laser is 1.5 W. At the end of evaporation we recompress the trap to the initial power to increase the thermalization rate and for a straightforward comparison of the NaLi density without re-scaling the trap volume. **a**, Single evaporation: (i) exponential forced evaporation ($\tau = 40$ ms) for 100 ms with trap depth $U(t) = A \exp(-t/\tau) + B$, where A and B are determined from the trap depths at $t = 0$ and 100 ms; (ii) recompression for 500 ms; (iii) hold for 40 ms for complete thermalization. **b**, Double evaporation: (1) exponential forced evaporation ($\tau = 40$ ms) for 100 ms; (2) recompression to the initial trap depth for 50 ms; (3) hold for 30 ms; (4) exponential evaporation ($\tau = 120$ ms) for 300 ms; (5) recompression for 200 ms.

200 ms, particles thermalize and the molecule temperature rises to the temperature of the heated Na atoms. When the trap amplitude is modulated in the same manner without the Na atoms, the temperature of molecules remains at 1.59(5) μ K. The heating process for sodium also induces centre-of-mass motion and breathing oscillations that cause the Na density to depend on time in the early stages of sympathetic heating, which is the reason for the delay in heating and for the non-exponential thermalization curve.

To measure the rate of thermalization, we return to the simpler situation of cooling in Fig. 2. We fit the temperature to a simple exponential model, $T(t) = (T_0 - T_\infty) \exp(-\Gamma_{\text{th}} t) + T_\infty$, where T_0 and T_∞ represent the initial and infinite-time-limit temperature of molecules, respectively, and obtain the thermalization rate, $\Gamma_{\text{th}} = 26(5) \text{ s}^{-1}$. Considering that thermalization requires about $3/\xi$ collisions (for Na–NaLi, $\xi \approx 1$; see Methods), we obtain the average elastic collision rate per particle, Γ_{el} , from the measured thermalization rate: $\Gamma_{\text{el}} \approx (3/\xi)\Gamma_{\text{th}} = 80(14) \text{ s}^{-1}$. In the presence of Na atoms, the initial loss rate of molecules is $\Gamma_{\text{inel}} = 1.29(6) \text{ s}^{-1}$, as obtained from a fit to the exponential loss model $N(t) = N_0 \exp(-\Gamma_{\text{inel}} t)$ (red dashed line in Fig. 2b). Comparing the average elastic collision rate to the total loss rate, we obtain the ratio of elastic to inelastic collisions for NaLi, $\gamma \approx \Gamma_{\text{el}}/\Gamma_{\text{inel}} = 62(12)$. Without Na atoms, the molecular loss follows a two-body loss model, $N(t) = N_0/(\beta t + 1)$ (blue solid line in Fig. 2b), from which we obtain an initial loss rate of $\beta = 1.02(6) \text{ s}^{-1}$. By considering the difference between Γ_{inel} and β as the effective inelastic loss rate for collisions between Na and NaLi, we obtain the ratio of ‘good’ to ‘bad’ collisions, $\gamma \approx \Gamma_{\text{el}}/(\Gamma_{\text{inel}} - \beta) \approx 300$ with an uncertainty of about 40%. The Na–NaLi loss rate constant is $4.0 \pm 1.3 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$; this is more than two orders of magnitude smaller than the universal loss model rate constant for Na–NaLi s-wave collisions³⁷, which is $1.7 \times 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ (see Methods). Whereas Na and NaLi in their upper stretched states form a stable mixture, as shown, preparing Na atoms in their lowest hyperfine state ($|F, m_F\rangle = |1, 1\rangle$) gives a loss rate consistent with the universal loss model³⁵.

We consider the two different evaporation ramps shown in Fig. 4. The molecule numbers, temperatures and PSDs resulting from these ramps are displayed in Fig. 5. For a single-species system with losses, optimal evaporation can be achieved by a single, continuous decrease of the trap depth. We achieve an increase in PSD by a factor of 7 by using a single stage of evaporation (Fig. 4a, red squares in Fig. 5). This increase is limited by the low initial Na density. In our system, thermalization is




Fig. 5 | Increasing phase-space density. **a–c**, The number **(a)**, temperature **(b)** and PSD **(c)** of NaLi molecules are plotted versus the power of the 1,596-nm trapping laser at the end of the forced exponential evaporation (i) for single evaporation and (1) for double evaporation in Fig. 4). For the double evaporation, we ramp the power down to 0.06 W (2 μ K trap depth) in (4), the lowest point that gives a molecular signal high enough for consistent temperature measurement. Single-stage evaporation to this low trap depth leads to loss of almost every molecule. The black dashed lines in **b** and **c** indicate the molecule temperature and PSD after STIRAP and removal of free atoms (shaded areas show ± 1 error bar). This PSD corresponds to $2.6(1) \times 10^4$ molecules. When Na atoms are not in the trap or when the evaporation is not performed to a sufficiently low trap depth, the PSD is lower than the initial value. This is mainly due to the molecular two-body loss. The solid curves are guides for the eye. Values and error bars are estimated as in Fig. 2 from 4–9 measurements.

dominated by the Na density, whereas loss is dominated by the NaLi density. To overcome the low Na density, we shorten the initial evaporation and recompression cycle ((1) and (2) in Fig. 4b). This cools the Na atoms and quickly increases the density of the cold Na for fast thermalization. After the cold, dense Na efficiently pre-cools the molecules during a short hold of 30 ms in the tight trap (3), we apply the second evaporation ramp (4). In this double evaporation, we achieve a peak PSD of $2.0(2) \times 10^{-2}$ (blue circles in Fig. 5), which is 20 times higher than the initial PSD. Before the final recompression of the trap, the lowest temperature of the molecules is 220(20) nK.

With Na present, more molecules survive the evaporation sequence as we evaporate Na atoms to a lower trap depth. This is mainly due to suppressed two-molecule loss. The loss rate constant scales linearly with the temperature owing to the p -wave character of the collisions³⁷. Evaporation to too low trap depth decreases the Na density, which makes the sympathetic cooling of molecules inefficient, and anti-evaporation (heating due to the preferential loss of the coldest molecules) dominates. This appears as a temperature increase below

a trap power of 0.17 W (5.7 μ K trap depth) for a single stage of evaporation, or 0.14 W (4.5 μ K trap depth) for two-stage evaporation. Without Na, the ramp-down in trap depth, (i), is adiabatic to a trap power of around 0.3 W, as evidenced by the constant molecule numbers and temperatures. Below that trap power, the ramp-down starts becoming non-adiabatic. As a result, a hot fraction of molecules escape, and both the molecule number and the temperature start decreasing. We note that because fermionic molecules do not thermalize by themselves, this temperature should be regarded as the average kinetic energy of molecules that are not in thermal equilibrium.

The favourable collisional properties of the spin-polarized Na–NaLi mixture in fully stretched hyperfine states result from strong suppression of electronic spin flips during collisions, which could otherwise lead to fast reactive or inelastic losses. In fully stretched states, direct spin exchange is forbidden and residual spin flips can occur only by weaker interactions. The main contribution to the weak spin flips is from dipolar relaxation due to direct coupling between the magnetic dipole moments of the spins of Na and NaLi. Studies in other spin-polarized systems^{17,34,38} predict spin flips induced by anisotropic electrostatic interaction at short range and intramolecular spin–spin and spin–rotation couplings to be weaker at ultracold temperatures. Dipolar relaxation can be eliminated by using the strong-field-seeking stretched state, but this was not necessary in our work. It is possible that the Na–NaLi system is favourable owing to the low reduced and molecular masses and the small relativistic effects, which result in a large rotational constant^{32,39}, low density of states²⁸ and small spin–orbit coupling⁴⁰, respectively. Further, ref.³¹ has shown that intramolecular spin–spin coupling causes spin flips, leading to chemical reactions for triplet NH molecules, thwarting evaporative cooling. Our results show a clear need for further work to determine what other molecules—and what conditions of internal states and external fields—would be suitable for collisional cooling at ultracold temperatures.

Technical upgrades to our apparatus should allow cooling into the quantum degenerate regime by increasing the initial Na density. The dominant loss mechanism—*p*-wave collisions between molecules—is expected to slow down compared to thermalization at lower temperatures, improving the efficiency of cooling. Further, the atomic and molecular states used in this work are magnetically trappable. Magnetic trapping would allow the use of much larger samples with perfect spin polarization. This would also eliminate the major concern of molecular loss induced by trapping light²⁹, making NaLi an ideal system for studying quantum chemistry. Concerns about sticky collisions²⁸, which can lead to a long-lived collision complex with high density of states, have led to pessimism over the prospects of achieving collisional cooling of ultracold molecules. This work demonstrates that sticky collisions do not limit NaLi, and probably other low-mass systems, suggesting a bright future for cooling light molecules.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-020-2141-z>.

- Anderson, M. H., Ensher, J. R., Matthews, M. R., Wieman, C. E. & Cornell, E. A. Observation of Bose–Einstein condensation in a dilute atomic vapor. *Science* **269**, 198–201 (1995).
- Davis, K. B. et al. Bose–Einstein condensation in a gas of sodium atoms. *Phys. Rev. Lett.* **75**, 3969–3973 (1995).
- Bloch, I., Dalibard, J. & Zwierger, W. Many-body physics with ultracold gases. *Rev. Mod. Phys.* **80**, 885–964 (2008).
- Baranov, M. A., Dalmonde, M., Pupillo, G. & Zoller, P. Condensed matter theory of dipolar quantum gases. *Chem. Rev.* **112**, 5012–5061 (2012).

- DeMille, D. Quantum computation with trapped polar molecules. *Phys. Rev. Lett.* **88**, 067901 (2002).
- Carr, L. D., DeMille, D., Krems, R. V. & Ye, J. Cold and ultracold molecules: science, technology and applications. *New J. Phys.* **11**, 055049 (2009).
- Christen, W., Rademann, K. & Even, U. Efficient cooling in supersonic jet expansions of supercritical fluids: CO and CO₂. *J. Chem. Phys.* **125**, 174307 (2006).
- Weinstein, J. D., deCarvalho, R., Guillet, T., Friedrich, B. & Doyle, J. M. Magnetic trapping of calcium monohydride molecules at millikelvin temperatures. *Nature* **395**, 148–150 (1998).
- Stellmer, S., Pasquiou, B., Grimm, R. & Schreck, F. Laser cooling to quantum degeneracy. *Phys. Rev. Lett.* **110**, 263003 (2013).
- Hu, J. et al. Creation of a Bose-condensed gas of ⁸⁷Rb by laser cooling. *Science* **358**, 1078–1080 (2017).
- Soldán, P. & Hutson, J. M. Interaction of NH(X² Σ^+) molecules with rubidium atoms: implications for sympathetic cooling and the formation of extremely polar molecules. *Phys. Rev. Lett.* **92**, 163202 (2004).
- Elioff, M. S., Valentini, J. J. & Chandler, D. W. Subkelvin cooling NO molecules via ‘billiard-like’ collisions with argon. *Science* **302**, 1940–1943 (2003).
- Henson, A. B., Gersten, S., Shagam, Y., Narevicius, J. & Narevicius, E. Observation of resonances in Penning ionization reactions at sub-kelvin temperatures in merged beams. *Science* **338**, 234–238 (2012).
- Tokunaga, S. K. et al. Prospects for sympathetic cooling of molecules in electrostatic, a.c. and microwave traps. *Eur. Phys. J. D* **65**, 141–149 (2011).
- González-Martínez, M. L. & Hutson, J. M. Ultracold hydrogen atoms: a versatile coolant to produce ultracold molecules. *Phys. Rev. Lett.* **111**, 203004 (2013).
- Lim, J., Frye, M. D., Hutson, J. M. & Tarbutt, M. R. Modeling sympathetic cooling of molecules by ultracold atoms. *Phys. Rev. A* **92**, 053419 (2015).
- Morita, M., Kosicki, M. B. Z., Żuchowski, P. S. & Tscherbul, T. V. Atom–molecule collisions, spin relaxation, and sympathetic cooling in an ultracold spin-polarized Rb(² Σ^+)–SrF(² Σ^+) mixture. *Phys. Rev. A* **98**, 042702 (2018).
- Augustovčová, L. D. & Bohn, J. Ultracold collisions of polyatomic molecules: CaOH. *New J. Phys.* **21**, 103022 (2019).
- Segev, Y. et al. Collisions between cold molecules in a superconducting magnetic trap. *Nature* **572**, 189–193 (2019).
- Reens, D., Wu, H., Langen, T. & Ye, J. Controlling spin flips of molecules in an electromagnetic trap. *Phys. Rev. A* **96**, 063420 (2017).
- Stuhl, B. K. et al. Evaporative cooling of the dipolar hydroxyl radical. *Nature* **492**, 396–400 (2012).
- Ni, K.-K. et al. A high phase-space-density gas of polar molecules. *Science* **322**, 231–235 (2008).
- Barry, J. F., McCarron, D. J., Norrgard, E. B., Steinecker, M. H. & DeMille, D. Magneto-optical trapping of a diatomic molecule. *Nature* **512**, 286 (2014).
- Anderegg, L. et al. Radio frequency magneto-optical trapping of CaF with high density. *Phys. Rev. Lett.* **119**, 103201 (2017).
- Truppe, S. et al. Molecules cooled below the doppler limit. *Nat. Phys.* **13**, 1173–1176 (2017).
- Collpy, A. L. et al. 3-D magneto-optical trap of yttrium monoxide. *Phys. Rev. Lett.* **121**, 213201 (2018).
- Żuchowski, Z. P. S. & Hutson, J. M. Reactions of ultracold alkali-metal dimers. *Phys. Rev. A* **81**, 060703 (2010).
- Mayle, M., Quémener, G., Ruzic, B. P. & Bohn, J. L. Scattering of ultracold molecules in the highly resonant regime. *Phys. Rev. A* **87**, 012709 (2013).
- Christiansen, A., Zwierlein, M. W., Groenenboom, G. C. & Karman, T. Trapping laser excitation during collisions limits the lifetime of ultracold molecules. *Phys. Rev. Lett.* **123**, 123402 (2019).
- Myatt, C. J., Burt, E. A., Ghrist, R. W., Cornell, E. A. & Wieman, C. E. Production of two overlapping Bose–Einstein condensates by sympathetic cooling. *Phys. Rev. Lett.* **78**, 586–589 (1997).
- Janssen, L. M. C., van der Avoird, A. & Groenenboom, G. C. Quantum reactive scattering of ultracold NH(X² Σ^+) radicals in a magnetic trap. *Phys. Rev. Lett.* **110**, 063201 (2013).
- Krems, R. V. & Dalgarno, A. Quantum-mechanical theory of atom–molecule and molecular collisions in a magnetic field: spin depolarization. *J. Chem. Phys.* **120**, 2296–2307 (2004).
- Żuchowski, P. S. & Hutson, J. M. Cold collisions of N(⁴S) atoms and NH(² Σ) molecules in magnetic fields. *Phys. Chem. Chem. Phys.* **13**, 3669–3680 (2011).
- Tscherbul, T. V., Ktos, J. & Buchachenko, A. A. Ultracold spin-polarized mixtures of ² Σ molecules with s-state atoms: collisional stability and implications for sympathetic cooling. *Phys. Rev. A* **84**, 040701 (2011).
- Rvachov, T. M. et al. Long-lived ultracold molecules with electric and magnetic dipole moments. *Phys. Rev. Lett.* **119**, 143001 (2017).
- Heo, M.-S. et al. Formation of ultracold fermionic NaLi Feshbach molecules. *Phys. Rev. A* **86**, 021602 (2012).
- Idziaszek, Z. & Julienne, P. S. Universal rate constants for reactive collisions of ultracold molecules. *Phys. Rev. Lett.* **104**, 113202 (2010).
- Hummon, M. T. et al. Cold N + NH collisions in a magnetic trap. *Phys. Rev. Lett.* **106**, 053201 (2011).
- Campbell, W. C. et al. Mechanism of collisional spin relaxation in ³ Σ molecules. *Phys. Rev. Lett.* **102**, 013003 (2009).
- Warehime, M. & Ktos, J. Nonadiabatic collisions of CaH with Li: importance of spin–orbit-induced spin relaxation in spin-polarized sympathetic cooling of CaH. *Phys. Rev. A* **92**, 032703 (2015).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2020

Methods

Atomic sample preparation

We produce an ultracold mixture of ^{23}Na and ^6Li in their upper stretched hyperfine states (in the $|F, m_F\rangle$ basis, these states correspond to $|2, 2\rangle$ for Na and $|3/2, 3/2\rangle$ for Li) in an Ioffe–Pritchard magnetic trap at a bias field of 2 G. We cool the Na atoms using microwave-induced evaporation and sympathetically cool the Li atoms with the Na atoms^{30,41–43}. Then, the mixture is transferred into the combination of two coaxial optical traps: a 1,064-nm optical dipole trap (30 μm waist, 10 W power) and a 1,596-nm 1D optical lattice (50 μm waist, 2 W power). After 0.6 s of forced evaporation in the 1,064-nm optical dipole trap, the trap is switched off. Then, in the 1,596-nm 1D lattice, the sample is transferred to the lowest-energy Zeeman states ($|1, 1\rangle$ for Na and $|1/2, 1/2\rangle$ for Li) using a Landau–Zener magnetic field sweep. By controlling the microwave power for the Landau–Zener sweep, we intentionally leave a fraction of Na atoms in the upper stretched hyperfine state for later use as coolants of the NaLi molecules.

Molecule formation and detection

As previously demonstrated with other atomic species^{22,35,44–51}, NaLi ground-state molecules are created from Feshbach molecules in the least-bound vibrational state. In a 1,596-nm 1D optical lattice, NaLi Feshbach molecules form in the $a^3\Sigma^+$, $\nu = 10$, $N = 2$, $m_N = -2$ state (ν and N are the vibrational and rotational molecular quantum numbers, respectively, and m_N is the magnetic quantum number of N) from the atoms in the lowest-energy Zeeman states by sweeping the magnetic field across an interspecies scattering resonance (Feshbach resonance) at 745 G (ref. ³⁶). Then, about 98% of Feshbach molecules are transferred to the triplet ground state ($a^3\Sigma^+$, $\nu = 0$, $N = 0$) by means of a 30- μs -long STIRAP sequence with an intermediate state ($\nu = 11$, $N = 1$, $m_N = -1$) in the $c^3\Sigma^+$ excited potential. The use of 1,596-nm light for the trap improves the lifetime of the NaLi molecules by suppressing spontaneous photon scattering compared to the more common 1,064-nm trap. The 1D lattice provides strong axial confinement to overcome the anti-trapping produced by magnetic curvature³⁵. The STIRAP light (upleg: 833 nm, 360.00381(1) THz; downleg: 819 nm, 366.23611(1) THz)^{35,52,53} is obtained from two home-built external cavity diode lasers (ECDL)⁵⁴ locked to an ultralow-expansion cavity. The relative linewidth of the two ECDLs is less than 1 kHz. To attain sufficient optical power for the upleg transition, we use a 500-mW high-power diode laser injection-locked by 2 mW of ECDL light. After molecule formation, a resonant light pulse is applied for about 1 ms to remove the free atoms that were not converted to molecules. These atoms in the lowest-energy Zeeman states accelerate the loss of NaLi to roughly the universal loss model rate³⁵. Sodium atoms in this state are optically pumped by the resonant removal light into a dark Zeeman state ($|F, m_F\rangle = |2, 1\rangle$). Adding a repump beam to address this state removes the need to reduce the magnetic field (by instantaneously shutting off the current in the coils generating the 745-G field), which caused considerable heating and loss of molecules in our previous work³⁵. We detect ground-state molecules using reverse STIRAP and a magnetic field sweep across the Feshbach resonance followed by resonant absorption imaging of the resulting free atoms.

Thermometry of molecules

To avoid issues with different atomic and molecular trapping potentials, we dissociate molecules in free space after turning off the trap, which allows accurate molecular thermometry. Given that the particle velocities are fixed after the trap is turned off, the density at some time of flight maps the velocity distribution of molecules in the trap. The molecule temperature is determined from a fit to such density profiles. Extra kinetic energy could be added in quadrature during the dissociation of Feshbach molecules into atomic continuum states⁵⁵. We determine the parameters of Feshbach dissociation (that is, the rate

of the magnetic field sweep) to minimize the released dissociation energy. All uncertainties quoted in this article are from statistical and fitting errors. The systematic uncertainty, which is mainly determined from the imaging magnification, is about 6% but has no effect on the measured thermalization rates because it gives a common multiplier to all temperature measurements.

Na–NaLi *s*-wave elastic-scattering cross-section

The elastic-scattering rate between Na and NaLi is given by $\Gamma_{\text{el}} = n_{\text{ov}}\sigma_{\text{el}}\nu_{\text{rel}}$ where n_{ov} is the overlap-averaged density of the mixture, σ_{el} is the elastic-scattering cross-section and ν_{rel} is the mean relative velocity⁵⁶:

$$\nu_{\text{rel}} = \sqrt{\frac{8k_B}{\pi} \left(\frac{T_{\text{Na}}}{m_{\text{Na}}} + \frac{T_{\text{NaLi}}}{m_{\text{NaLi}}} \right)}$$

$$n_{\text{ov}} = \frac{I}{\frac{N_{\text{Na}}N_{\text{NaLi}}}{N_{\text{Na}} + N_{\text{NaLi}}}} = \frac{N_{\text{Na}}N_{\text{NaLi}}\bar{\omega}_{\text{Na}}^3 \left[\frac{2\pi k_B T_{\text{Na}}}{m_{\text{Na}}} \left(1 + \frac{\alpha_{\text{Na}}}{\alpha_{\text{NaLi}}} \frac{T_{\text{NaLi}}}{T_{\text{Na}}} \right) \right]^{-3/2}}{\frac{N_{\text{Na}}N_{\text{NaLi}}}{N_{\text{Na}} + N_{\text{NaLi}}}} \quad (1)$$

where m_i , T_i and α_i represent the mass, temperature and polarizability, respectively, of particle i , $I = \int dV n_{\text{Na}} n_{\text{NaLi}}$ is the density overlap integral of the mixture in a harmonic trap, $\bar{\omega}_{\text{Na}} = (\omega_x \omega_y \omega_z)^{1/3} = 2\pi \times (540 \times 410 \times 34,000)^{1/3}$ Hz is the geometric mean of the trap frequencies, and k_B is the Boltzmann constant. The differential gravitational sag between molecules and atoms has a negligible effect on the overlap density. The effective number of particles per lattice site, $N_{(\text{Na}, \text{NaLi})}$, is the total number of particles divided by the factor l_{eff}/a , where $a = \lambda/2$ is the lattice spacing (λ , wavelength of the trap laser) and l_{eff} is the effective axial width (in the z direction; see Fig. 1) of the atomic density distribution across all lattice sites:

$$l_{\text{eff}} = 2 \int_0^\infty z \exp(-z^2/2\sigma^2) dz = \sqrt{\frac{8}{\pi}} \sigma \quad (2)$$

where $\sigma = 0.72(1)$ mm is the Gaussian width, obtained from a fit of the atomic density profile.

In a mass-imbalanced system, the factor $3/\xi$ quantifies the approximate average number of collisions per particle required for thermalization, where $\xi = 4m_{\text{Na}}m_{\text{NaLi}}/(m_{\text{Na}} + m_{\text{NaLi}})^2$ (ref. ⁴²). For the Na–NaLi mixture, this corresponds to approximately three collisions because the mass difference is small. Thus, the relation between the thermalization rate and the elastic-scattering rate is given by $\Gamma_{\text{th}} \approx \Gamma_{\text{el}}/(3/\xi)$. In our system, where the particle number is largely imbalanced (that is, $N_{\text{Na}}/N_{\text{NaLi}} \approx 7$), we can write the thermalization rate as:

$$\Gamma_{\text{th}} \approx \frac{(I/N_{\text{NaLi}})\sigma_{\text{el}}\nu_{\text{rel}}}{3/\xi} \quad (3)$$

where I/N_{NaLi} is the average density of Na atoms seen by the NaLi molecules. Given the measured thermalization rate, we obtain the elastic-scattering cross-section between a molecule and an atom, $\sigma_{\text{el}} \approx (3/\xi)(N_{\text{NaLi}}/\nu_{\text{rel}})I\Gamma_{\text{th}} = 2.4(5) \times 10^{-11}$ cm², and the corresponding scattering length, $a = 263(24)a_0$, where a_0 is the Bohr radius.

In addition to the simple exponential fit, we perform a numerical simulation of the thermalization process based on the differential equation for $\Delta T = T_{\text{NaLi}} - T_{\text{Na}}$ (refs. ^{42,43,56}):

$$\frac{1}{\Delta T} \frac{d(\Delta T)}{dt} = -\Gamma_{\text{th}} = -\frac{\xi}{3} n_{\text{ov}} \sigma_{\text{el}} \nu_{\text{rel}} \quad (4)$$

In the numerical simulation, the temperature dependence of n_{ov} , σ_{el} and ν_{rel} is fully taken into account. The *s*-wave elastic-scattering cross-section between Na and NaLi is given by $\sigma_{\text{el}} = \frac{4\pi a^2}{[1 - (1/2)k^2 r_0 a]^2 + (ka)^2}$, where

Article

$\hbar k = \mu v_{\text{rel}}$ is the relative momentum (μ is the reduced mass, \hbar is the reduced Planck constant) and r_0 is the effective range calculated from the scattering length, a , and from $C_6^{\text{Na-NaLi}}$ (see Methods section ‘Na–NaLi loss rate constant from the universal loss model’). We iteratively solve equation (4) by varying a , and find the optimal solution that minimizes the weighted sum of squared residuals: $\sum_i [(\Delta T_{m,i} - \Delta T_{n,i})/\sigma_{\Delta T_{m,i}}]^2$, where the subscript n denotes the numerical solution, m denotes the measurement and $\sigma_{\Delta T_{m,i}}$ is the uncertainty of the measurement at the i th time step, estimated as in Fig. 2. The numerical solution for a obtained in this way agrees with the result calculated from the simple exponential model described above within the uncertainty.

Na–NaLi loss rate constant

The Na–NaLi loss rate constant is defined as the differential loss rate, $\Gamma_{\text{inel}} - \beta$, normalized by the density of Na overlapped with NaLi (that is, the overlap-averaged density of the mixture, n_{ov}), where Γ_{inel} is the molecular loss rate with Na and β is without Na. As shown in the calculation of the s -wave elastic-scattering cross-section between Na and NaLi, $n_{\text{ov}} \approx I/N_{\text{NaLi}}$, given the large imbalance of particle numbers. Thus, the Na–NaLi loss rate constant mentioned in the main text is calculated from $(\Gamma_{\text{inel}} - \beta)/(I/N_{\text{NaLi}})$.

Na–NaLi loss rate constant from the universal loss model

We compare the measured loss rate constant for Na and NaLi with the theoretical prediction based on the universal loss model³⁷. When calculating the theoretical loss rate constant, we estimate the van der Waals coefficient between Na and NaLi, $C_6^{\text{Na-NaLi}}$, as the sum of the C_6 coefficients for the atomic pairs that constitute the collision partners, using values from ref. 57: $C_6^{\text{Na-NaLi}} = C_6^{\text{Na-Li}} + C_6^{\text{Na-Na}}$.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

41. Hadzibabic, Z. et al. Fiftyfold improvement in the number of quantum degenerate fermionic atoms. *Phys. Rev. Lett.* **91**, 160401 (2003).

42. Mosk, A. et al. Mixture of ultracold lithium and cesium atoms in an optical dipole trap. *Appl. Phys. B* **73**, 791–799 (2001).
43. Ivanov, V. V. et al. Sympathetic cooling in an optically trapped mixture of alkali and spin-singlet atoms. *Phys. Rev. Lett.* **106**, 153201 (2011).
44. Lang, F., Winkler, K., Strauss, C., Grimm, R. & Denschlag, J. H. Ultracold triplet molecules in the rovibrational ground state. *Phys. Rev. Lett.* **101**, 133005 (2008).
45. Danzl, J. G. et al. An ultracold high-density sample of rovibronic ground-state molecules in an optical lattice. *Nat. Phys.* **6**, 265 (2010).
46. Takekoshi, T. et al. Ultracold dense samples of dipolar RbCs molecules in the rovibrational and hyperfine ground state. *Phys. Rev. Lett.* **113**, 205301 (2014).
47. Molony, P. K. et al. Creation of ultracold ⁸⁷Rb¹³³Cs molecules in the rovibrational ground state. *Phys. Rev. Lett.* **113**, 255301 (2014).
48. Park, J. W., Will, S. A. & Zwierlein, M. W. Ultracold dipolar gas of fermionic ²³Na⁴⁰K molecules in their absolute ground state. *Phys. Rev. Lett.* **114**, 205302 (2015).
49. Guo, M. et al. Creation of an ultracold gas of ground-state dipolar ²³Na⁸⁷Rb molecules. *Phys. Rev. Lett.* **116**, 205303 (2016).
50. Yang, H. et al. Observation of magnetically tunable Feshbach resonances in ultracold ²³Na⁴⁰K + ⁴⁰K collisions. *Science* **363**, 261–264 (2019).
51. Seeßelberg, F. et al. Extending rotational coherence of interacting polar molecules in a spin-decoupled magic trap. *Phys. Rev. Lett.* **121**, 253401 (2018).
52. Rvachov, T. M. et al. Photoassociation of ultracold NaLi. *Phys. Chem. Chem. Phys.* **20**, 4746–4751 (2018).
53. Rvachov, T. M. et al. Two-photon spectroscopy of the NaLi triplet ground state. *Phys. Chem. Chem. Phys.* **20**, 4739–4745 (2018).
54. Cook, E. C., Martin, P. J., Brown-Heft, T. L., Garman, J. C. & Steck, D. A. High passive-stability diode-laser design for use in atomic-physics experiments. *Rev. Sci. Instrum.* **83**, 043101 (2012).
55. Mukaiyama, T., Abo-Shaer, J. R., Xu, K., Chin, J. K. & Ketterle, W. Dissociation and decay of ultracold sodium molecules. *Phys. Rev. Lett.* **92**, 180402 (2004).
56. Ravensbergen, C. et al. Production of a degenerate Fermi–Fermi mixture of dysprosium and potassium atoms. *Phys. Rev. A* **98**, 063624 (2018).
57. Derevianko, A., Babb, J. F. & Dalgarno, A. High-precision calculations of van der Waals coefficients for heteronuclear alkali-metal dimers. *Phys. Rev. A* **63**, 052704 (2001).

Acknowledgements We thank M. Zwierlein for discussions and J. Yao for technical assistance. We acknowledge support from the NSF through the Center for Ultracold Atoms and award 1506369, from the NASA Fundamental Physics Program and from the Samsung Scholarship.

Author contributions H.S., W.K. and A.O.J. conceived the experiment. H.S. led the data measurement. H.S. and A.O.J. performed the data analysis. H.S., J.J.P. and A.O.J. designed and constructed the experimental setup. All authors discussed the results and contributed to the writing of the manuscript.

Competing interests The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to H.S.

Reprints and permissions information is available at <http://www.nature.com/reprints>.